REPORT DOCUMENTATION PAGE					Form Approved OMB No. 0704-0188	
data needed, and completing a this burden to Department of D 4302. Respondents should be	and reviewing this collection of in refense, Washington Headquart aware that notwithstanding any	nformation. Send comments rega ers Services, Directorate for Infor tother provision of law, no persor	arding this burden estimate or an mation Operations and Reports a shall be subject to any penalty	y other aspect of this co (0704-0188), 1215 Jeffe	thing existing data sources, gathering and maintaining the illection of information, including suggestions for reducing erson Davis Highway, Suite 1204, Arlington, VA 22202-11 a collection of information if it does not display a currently	
1. REPORT DATE (DD		R FORM TO THE ABOVE ADDE	RESS.	3. D	DATES COVERED (From - To)	
05/30/2012	,	Final Report			9/01/10 - 02/29/12	
4. TITLE AND SUBTIT		-			CONTRACT NUMBER	
Plasmonic-Enha	nced Catalysis	5		FA	9550-10-1-0504	
				5b.	GRANT NUMBER	
					PROGRAM ELEMENT NUMBER	
				MO	D: P00001	
6. AUTHOR(S)				5d.	PROJECT NUMBER	
Alivisatos, A. Paul,	Somorjai, Gabor A.,	and Yang, Peidong				
				5e.	TASK NUMBER	
				5f. \	WORK UNIT NUMBER	
7 PERFORMING ORG	SANIZATION NAME(S)	AND ADDRESS(ES)		8 P	ERFORMING ORGANIZATION REPORT	
Regents of the University of California The University of California, Berkeley				N	### 1000 PART PART	
Regents of the University of				02	9913-003: APAZ01ZAFOSR	
California	e onitversity of	-				
	Avenue, Rm. 31	3				
Berkeley, CA	•					
9. SPONSORING / MONITORING AGENCY NAME(S) AND ADDRESS(ES)					SPONSOR/MONITOR'S ACRONYM(S)	
AFOSR/PKR2				003	2416 DAF AFOSR	
USAF, AFRL, DU						
AF Ofc of Scientific Research					SPONSOR/MONITOR'S REPORT	
875 Randolph St, Rm 3112					NUMBER(S)	
Arlington, VA 22203 12. DISTRIBUTION / AVAILABILITY STATEMENT				Al	FRL-OSR-VA-TR-2012-1020	
	public distrik					
13. SUPPLEMENTAR	Y NOTES					
hot electron flow we an iron oxide photoa catalytic intermediat associated with the l nanoparticles that we surface enhanced rai	re explored. Light transcribed by more than 5 es were demonstrated ocalized surface plas ould be lost by ensen man spectroscopy (SI	apping from nanostruction of across the solar spans. Firstly, single particular mon resonance (LSPR) able averaging were of ERS) measurements were supported to the state of t	etured Au nanopillars bectrum. Two platform the hydrogen storage to the hydrogen storage to the hydrogen storage to the hydrogen storage. Shapes because the hydrogen storage to the hydrogen storage	and their surfactors for in situ plate rajectories were ape dependent have Ag-alumina late totooxidation of	catalysis and surface plasmon driven e plasmon enhanced the photocurrent of smon enhanced spectroscopy of e monitored by recording the scattering sydrogen storage trajectories of Pd hybrid platforms for reproducible f water on titiania was shown to occur are spectroscopically silent or weak to	
45 OUD 1507 757						
15. SUBJECT TERMS plasmonic cata		en storage, SER	S, single part:	icle trajec	tories	
16. SECURITY CLASSIFICATION OF:			17. LIMITATION	18. NUMBER	19a. NAME OF RESPONSIBLE PERSON	
			OF ABSTRACT	OF PAGES	A. Paul Alivisatos	
a. REPORT	b. ABSTRACT	c. THIS PAGE			19b. TELEPHONE NUMBER (include area code)	

510-486-5111

Comprehensive summary of significant work:

- Researchers followed the trajectories of hydrogen storage in single Pd nanoparticles using Au as a plasmonic probe, with
 direct and indirect sensing schemes. Hydrogen absorption and desorption trajectories from individual particles were
 disentangled from ensemble averaging.
- In indirect sensing scheme, a Pd dot is patterned in the near-field of an Au antenna with great precision. Geometric enhancement of the electromagnetic field with triangular Au antennas results in a larger sensing response compared to Au rod antennas.
- Au/Pd core/shell nanoparticles for the direct sensing experiment yielded larger LSPR shifts as both materials were in contact. Notably, shape dependent hydrogen uptake trajectories were observed. The majority of these core-shell particles displayed red and blue LSPR shifts with increasing and decreasing partial pressures of hydrogen respectively, as hydrogen intercalated reversibly into palladium. However, 20% of the particles, mainly those with thicker Pd shells, showed LSPR shifts that cannot be merely explained hydride formation. These unexpected LSPR shifts may be explained by Au/Pd interdiffusion or silicide formation.
- This non-invasive, in situ sensing platform underlies the need for more single particle studies of catalytically active nanoparticles. This work shows that single particle techniques are vital in retaining information that would be lost in ensemble averaging.
- Hot electron flow generated on a gold thin film by photon absorption (or internal photoemission) was amplified by localized surface plasmon resonance.
- Direct measurement of photocurrent on a chemically modified gold thin film of metal-semiconductor (TiO₂) Schottky diodes
- Intrinsic correlation between the hot electron flow generated by internal photoemission and localized surface plasmon.
- This project demonstrated enhanced photocurrent in a thin-film iron oxide photoanode coated on arrays of Au nanopillars. A net enhancement as high as 50% was observed over the solar spectrum. The enhancement can be attributed primarily to the increased optical absorption originating from both surface plasmon resonances and photonic-mode light trapping in the nanostructured topography.
- Project demonstrated a new type of Ag-alumina hybrid SERS platforms for detection of surface reaction spectroscopically in real time. The reaction intermediates of water oxidation on TiO₂ surface were observed in real time under UV irradiation. Surface intermediate bands, which correspond to peroxo, hydroperoxo and hydroxo of titania surface intermediate species were clearly observed, suggesting that the photooxidation of water splitting was initiated by nucleophilic attack mechanism.

Catalytically active nanoparticles were interrogated at the single particle level using a plasmonic sensing scheme. The activity and selectivity of heterogenous catalysts is uniquely dependent on their size, shape, faceting and support. Heterogenous catalyst particles also tend to evolve as the reaction proceeds. Single particle studies of catalytically active nanoparticles deconvolute the trajectory of individual particles, retaining information that would be lost by ensemble averaging. Hydrogen storage was chosen as a model reaction to study catalyst nanoparticles at the single particle level. As hydrogen gas dissociates and intercalates into Pd, Pd changes from a metal to a semi-metal. This change in the dielectric function of Pd gives rise to a change in the refractive index in the near-field of the Au antenna, shifting its localized surface plasmon resonance (LSPR). More oblate particles show a larger sensing response due to field enhancement. Shape dependent hydrogen storage trajectories are observed for Au/Pd core/shell nanocrystals.

Our second plasmon enhanced sensing scheme of catalytically active intermediates involved a new type of Ag-alumina hybrid SERS platform for spectroscopy of surface reactions in real time. Langmuir-Blodgett technique was employed as a large scale assembly technique to self-assemble facetted Ag nanocrystals into monolayers of sensor arrays with reproducible SERS signals. The Ag nanostructures platforms were then coated with a thin layer of alumina by atomic layer deposition to prevent charge transfer between photocatalyst and Ag during photocatalysis. A single layer of ~ 5 nm thick Pt co-catalyst coated anatase TiO₂ nanocrystals with predominantly $\{001\}$ facets were subsequently assembled onto the SERS platforms by Langmuir-Blodgett technique. The surface-enhanced Raman vibrations of distorted surface Ti-O-Ti bands and reaction intermediates of water oxidation on TiO₂ surface were observed in real time under UV irradiation. Surface intermediate bands, which correspond to peroxo, hydroperoxo and hydroxo of titania surface intermediate species were clearly observed. The detection of titania peroxo and hydroperoxo species suggested the photooxidation of water splitting was initiated by nucleophilic attack mechanism.

In summary, researchers developed two plasmon enhanced sensing platforms for the in situ detection of intermediates in catalysis. They also examined the plasmonic enhancement of iron oxide photoanodes, and the amplification of hot electrons by the localized surface plasmon resonance of gold. This body of work illuminated spectroscopically silent or weak intermediates, and thus contributed directly to the rational design of catalysts.

Papers published:

"Surface Plasmon-Driven Hot Electron Flow Probed with Metal-semiconductor Nanodiodes," Y. K. Lee, C. H. Jung, J. Park, H. Seo, G. A. Somorjai, J. Park. Nano Lett. 11, 4251, 2011.

"Templated Synthesis of Shape Controlled Ordered TiO2 Cage Structures," Y. Deng, H. Tüysüz, J. Henzie, P. Yang, Small, 7, 2037, 2011.

"Observations of shape-dependent Hydrogen Uptake Trajectories from Single Nanocrystals," M. L. Tang, N. Liu, J. A. Dionne, A. P. Alivisatos, J. Am. Chem. Soc. 133, 13220, 2011.

"Nanoantenna-enhanced Gas Sensing in a Single Tailored Nanofocus," N. Liu, M. L. Tang, M. Hentschel, H. Giessen, A. P. Alivisatos, Nature Mater. 10, 631, 2011. (This was selected as the cover for the August 2011 issue.)

"Plasmon Enhanced-Photocatalytic Activity of Iron Oxide on Gold Nanopillars," H. Gao, C. Liu, H. E. Jeong, P. Yang, ACS Nano. 6, 234, 2012.

"Ag Nanoparticle-Alumina Hybrid Surface-Enhanced Raman Spectroscopy Platforms for Direct Spectroscopic Detection of Overall Water Splitting Reaction Intermediates," X. Y. Ling, S. Lo, D. Hoang, C. Liu, M. Fardy, P. Yang, J. Am. Chem. Soc. (in preparation).

Project Staff:

PIs: A. Paul Alivisatos, Gabor A. Somorjai, Peidong Yang Postdoctoral Scholars and Graduate Students: Yimin Lee, Na Liu, Ming Lee Tang, Rouxie Yan, Jesse Engel, David Grauer